arXiv:1604.04807v2 [math-ph] 27 Nov 2016

On some integrable Emden-Fowler equations which yield Weierstrass elliptic solutions

Stefan C. Mancas* and Haret C. Rosu†

*Department of Mathematics, Embry–Riddle Aeronautical University
Daytona Beach, FL. 32114-3900, U.S.A.

†IPICyT, Instituto Potosino de Investigacion Cientifica y Tecnologica,
Camino a la presa San José 2055, Col. Lomas 4a Sección,
San Luis Potosí, C.P. 78216, S.L.P., Mexico

Abstract

We discuss two integrable classes of the Emden-Fowler (EF) equation of the type $z'' = A\chi^{-\lambda-2}z^n$ for $\lambda = \frac{n-1}{2}$, and $\lambda = n+1$, by finding their corresponding invariants which will reduce them to first order nonlinear ordinary differential equations (ODEs). By using particular solutions of the EF equations, we reduce both classes to the form $\ddot{\nu} + a\dot{\nu} + b(\nu - \nu^n) = 0$ where a, b depend only on λ, n, and for both cases we write closed form parametric solutions. The illustrative examples are taken from astrophysics and general relativity and correspond to two $n = 2$ cases from class one and two, and one $n = 5$ case from class one, all of them yielding Weierstrass elliptic solutions.

PACS numbers: 02.30.Gp, 02.30.Hq, 02.30.Ik, 02.30.Jr

Keywords: Emden-Fowler equation, Lane-Emden equation, elliptic functions, Weierstrass functions, parametric solutions.

1 Introduction

Several particular cases of the homogeneous ODEs of the form

$$y''(x) + \frac{N}{x}y'(x) + f(x,y) = 0 \ , \tag{1}$$

where $N > 0$, $x \geq 0$, and $f(x,y)$, a nonlinear function, have proved to be extremely significant in fundamental topics of astrophysics, atomic physics, and other areas. The first occurrence was in astrophysics in the particular case $N = 2$ and $f(x,y)$ a monomial power of y

$$y''(x) + \frac{2}{x}y'(x) + y^n = 0 \ , \tag{2}$$

or, in self-adjoint form

$$\frac{d}{dx}\left(x^2\frac{dy}{dx}\right) + x^2y^n = 0 \ , \tag{3}$$

with $n = 3$ the most natural case, which is known as the Lane-Emden equation. For standard initial conditions $y(0) = 1$ and $y'(0) = 0$, there are solutions of Eq. (2) representing the Newton-Poisson gravitational

*E-mail address: mancass@erau.edu
†E-mail address: hcr@ipicyt.edu.mx

potential of stars, such as the Sun, considered as spheres filled with polytropic gas. This has been first shown in the famous book of Emden [1] written at the beginning of the 20th century. Emden's book not only summarized the first four decades of research on the self-gravitating stars initiated by Lane in 1870 [2], but also introduced one of the first singular Cauchy problem as represented by Eq. (2), and moreover stimulated further remarkable studies of a whole generation of renowned astrophysicists and mathematicians, such as Eddington [3], Fowler [4], and Milne [5]. Furthermore, for negative values of the parameter n there are important applications in the theory of spherical nebulae and clusters [6] and in the Emden-Fowler form to more terrestrial areas such as pseudoplastic fluids and large deflections of membranes in mechanics [7].

A 'two-parameter' generalization of the Emden equation was introduced and systematically studied by Fowler [4]

$$\frac{d}{dx}\left(x^2\frac{dy}{dx}\right)+x^{\lambda}y^n=0 \tag{4}$$

and is currently known as the Emden-Fowler equation, while the Lane-Emden equation is the particular case $\lambda=2$. By a change of variables, the EF equation can be written in the standard, non-adjoint form

$$\frac{d^2z}{d\chi^2}+\chi^{-\lambda-2}z^n=0\ , \tag{5}$$

which is the usual form encountered in the literature.

As for the 'three-parameter' generalization

$$\frac{d}{dx}\left(x^{\rho}\frac{dy}{dx}\right)\pm x^{\lambda}y^n=0 \tag{6}$$

a similar change of variable reduces it to the standard form

$$\frac{d^2z}{d\chi^2}\pm\chi^{\sigma}z^n=0\ , \tag{7}$$

which is equivalent to Eq. (5) and so nothing essentially new is added. The Thomas-Fermi model [8, 9] for the electrostatic field in the bulk of a heavy atom originates from Coulomb-Poisson considerations and is expressed by a self-adjoint form of the kind given in Eq. (6) although for a non-integer n.

In this work, we are concerned with the standard form of the EF equation in the form

$$\frac{d^2z}{d\chi^2}=A\chi^{-\lambda-2}z^n\ , \tag{8}$$

for which we provide a discussion of two important integrable cases corresponding to $\lambda=\frac{n-1}{2}$ and $\lambda=n+1$.

There are two particular cases of Eq. (8). First, for $\lambda=0$ we obtain

$$\frac{d^2z}{d\chi^2}=A\chi^{-2}z^n\ , \tag{9}$$

which is transformed to

$$\ddot{z}-\dot{z}=Az^n \tag{10}$$

by using $t=\ln\chi$. The second particular case is $\lambda=n-1$ which leads to

$$\frac{d^2z}{d\chi^2}=A\chi^{-n-1}z^n\ , \tag{11}$$

and using again $t=\ln\chi$ simplifies to

$$\ddot{z}-\dot{z}=A\chi^{-n+1}z^n\ . \tag{12}$$

Moreover, by letting $z=\theta e^t$, we obtain

$$\ddot{\theta}+\dot{\theta}=A\theta^n\ . \tag{13}$$

Equations (10), (13) were discussed previously in detail by [10, 4] and will not be of interest here.

2 Self-adjoint and invariant forms

The self-adjoint form of Eq. (8) is obtained by changing the variables according to $z(\chi) = \eta(\xi)$ and $\xi = \frac{1}{\chi}$, which leads to

$$\frac{1}{\xi^2}\frac{d}{d\xi}(\xi^2\eta_\xi) = A\xi^{\lambda-2}\eta^n \ . \tag{14}$$

On the other hand, an invariant form of Eq. (8) can be obtained using the transformation $z(\chi) = \frac{w(s)}{s}$, where $s = \frac{1}{\chi}$, which leads to an EF equation with a different power of the independent variable containing both parameters

$$w_{ss} = As^{\lambda-1-n}w^n. \tag{15}$$

One can then infer that if $\lambda = \frac{n-1}{2}$ Eq. (15) is invariant under the transformation $z \leftrightarrow \frac{w(s)}{s},\ s \leftrightarrow \frac{1}{\chi}$ since it coincides with Eq. (8), while for $\lambda = n+1$, the power of s is zero. Thus, it does not appear to be a coincidence that in 1984 when Rosenau [12] was interested in the integration of Eq. (8) where he provided two algebraic methods of integration when the same conditions between the powers of the variables are satisfied: $\lambda = \frac{n-1}{2}$ or $\lambda = n+1$. Thus, both cases will be considered in detail for which we will construct integrals of motions and write the parametric solutions using a general formalism.

To proceed with the general solutions for these two classes first let us assume we have a particular solution $z_p(\chi)$ of Eq. (8), then we construct a general solution $z(\chi) = z_p(\chi)\nu(\chi)$, where $\nu(\chi)$ must satisfy the ODE

$$z_p\nu'' + 2z_p{}'\nu' + z_p{}''(\nu - \nu^n) = 0. \tag{16}$$

For both classes the coefficients $\frac{z_p{}'}{z_p}$ and $\frac{z_p{}''}{z_p}$ turn to be constants.

2.1 Case $\lambda = \frac{n-1}{2}$

Let $\lambda = \frac{n-1}{2}$ in Eq. (8) we obtain

$$z'' = A\chi^{-\frac{n+3}{2}}z^n \ , \tag{17}$$

which has particular solution

$$z_p(\chi) = \alpha\sqrt{\chi}\,, \qquad \alpha^{n-1} = -\frac{1}{4A}. \tag{18}$$

To construct the integrals of motion we first put Eq. (17) in self-adjoint form to obtain

$$\frac{d}{d\xi}(\xi^2\eta_\xi) = A\xi^{\frac{n-1}{2}}\eta^n \ . \tag{19}$$

In 1970s, Djukic [13] was first to find the integrals of motion for Eq. (17) using a generalized Noether theorem. He obtained

$$\xi^2\eta_\xi(\xi\eta_\xi + \eta) - \frac{2A}{n+1}\xi^{\frac{n+1}{2}}\eta^{n+1} = \mathcal{C}_1, \tag{20}$$

which in terms of the original variables of Eq. (17) turns into

$$z'(\chi z' - z) - \frac{2A}{n+1}\chi^{-\frac{n+1}{2}}\chi^{n+1} = \mathcal{C}_1. \tag{21}$$

The invariant method may have advantage over the original equation because instead of solving a second order ODE, one may obtain a solution by solving the first order ODE first.

Using the invariant form (15) with $\lambda = \frac{n-1}{2}$ we obtain

$$w_{ss} = As^{-\frac{n+3}{2}}w^n \ , \tag{22}$$

which is the same equation as (17). Thus, its invariant is

$$w_s(sw_s - w) - \frac{2A}{n+1} s^{-\frac{n+1}{2}} w^{n+1} = \mathcal{C}_2 \ . \tag{23}$$

By using the original variables, the invariant (23) is exactly (21) up to some arbitrary integration constant.

Since we know z_p according to (18) then Eq. (16) reduces to

$$\chi^2 \nu'' + \chi \nu' - \frac{1}{4}(\nu - \nu^n) = 0 \ . \tag{24}$$

The damping term can be eliminated using $t = \ln \chi$ and gives

$$\ddot{\nu} = \frac{1}{4}(\nu - \nu^n) \ , \tag{25}$$

with solution found by Fowler [14]

$$\nu(t) = \left[2K_1(n+1) \frac{e^{-\frac{n-1}{2}t}}{\left(1 + K_1 e^{-\frac{n-1}{2}t}\right)^2} \right]^{\frac{1}{n-1}} \ , \qquad n \neq 1 \ . \tag{26}$$

By combining Eqs. (26) and (18), the solution to Eq. (17) is

$$z(\chi) = \left[\frac{-K_1(n+1)}{2A\left(1 + K_1 \chi^{-\frac{n-1}{2}}\right)^2} \right]^{\frac{1}{n-1}} \ , \qquad n \neq 1 \ . \tag{27}$$

The case $n = 1 \Rightarrow \lambda = 0$ gives Eq. (10).

2.2 Case $\lambda = n + 1$

Letting $\lambda = n + 1$ in Eq. (8) we get

$$z'' = A\chi^{-(n+3)} z^n \ , \tag{28}$$

which has particular solution

$$z_p(\chi) = \beta \chi^{\frac{n+1}{n-1}} \ , \qquad \beta^{n-1} = \frac{2(n+1)}{A(n-1)^2} \ , \qquad n \neq 1. \tag{29}$$

To construct the integrals of motion for the above equation we will proceed in reversed order, by using first the invariant transformation and then the canonical variables. Using $\lambda = n + 1$ in Eq. (15) we have

$$w_{ss} = A w^n \ . \tag{30}$$

The integral of motion is easily obtained by multiplying by w_s and integrating to get

$$w_s^2 - \frac{2A}{n+1} w^{n+1} = \mathcal{C}_3 \tag{31}$$

which in terms of the original variables of Eq. (28) becomes

$$(z - \chi z')^2 - \frac{2A}{n+1} \chi^{-(n+1)} z^{n+1} = \mathcal{C}_3 \ . \tag{32}$$

Using canonical variables, Eq. (28) is

$$\frac{d}{d\xi}(\xi^2 \eta_\xi) = A \xi^{n+1} \eta^n \ . \tag{33}$$

Therefore, by using the canonical variables in Eq. (32) we obtain the invariant

$$(\eta+\xi\eta_\xi)^2-\frac{2A}{n+1}\xi^{n+1}\eta^{n+1}=\mathcal{C}_3\ . \tag{34}$$

Since we know z_p, then according to Eq. (16) we obtain

$$\chi^2v''+\frac{2(n+1)}{n-1}\chi v'+\frac{2(n+1)}{(n-1)^2}(v-v^n)=0\ . \tag{35}$$

Using $t=\ln\chi$ we obtain

$$\ddot{v}+\frac{n+3}{n-1}\dot{v}+\frac{2(n+1)}{(n-1)^2}(v-v^n)=0\ . \tag{36}$$

A particular solution of this equation can be obtained using the factorization technique of Rosu and Cornejo-Pérez [15]. The factored form of Eq. (35) is

$$\left[\frac{d}{dt}+\frac{n+1}{n-1}\left(1+\nu^{\frac{n-1}{2}}\right)\right]\left[\frac{d}{dt}+\frac{2}{n-1}\left(1-\nu^{\frac{n-1}{2}}\right)\right]\nu=0\ . \tag{37}$$

This leads to

$$\dot{\nu}+\frac{2}{n-1}\nu-\frac{2}{n-1}\nu^{\frac{n+1}{2}}=0 \tag{38}$$

which is a Bernoulli equation with kink solution

$$\nu(t)=\frac{1}{(1+K_2e^t)^{\frac{2}{n-1}}}\ ,\qquad n\neq1\ . \tag{39}$$

By combining Eqs. (39) and (29), the solution to Eq. (28) is

$$z(\chi)=\left[\frac{2(n+1)}{A(n-1)^2}\frac{\chi^{n+1}}{(1+K_2\chi)^2}\right]^{\frac{1}{n-1}}\ ,\qquad n\neq1\ . \tag{40}$$

3 Parametric solutions

3.1 Case $\lambda=\frac{n-1}{2}$

Eq. (17) is Eq. (4) in section §2.3.1-2 of Polyanin's book [16]. Depending on n there are two sets of parametric solutions,

$$\begin{aligned}
&\chi(\tau)=aB_1^2\exp\Theta(\tau)\\
&z(\tau)=bB_1\tau\exp\tfrac{1}{2}\Theta(\tau)\\
&\Theta(\tau)=\int\frac{d\tau}{\sqrt{K_3+\psi(\tau)+\frac{\tau^2}{4}}}\\
&A=\left(\tfrac{a}{b^2}\right)^{\frac{n-1}{2}}\ .
\end{aligned} \tag{41}$$

First, if $n\neq-1$ we have $\psi(\tau)=\frac{2\tau^{n+1}}{n+1}$, while if $n=-1$ then $\psi(\tau)=2\ln|\tau|$.

3.2 Case $\lambda=n+1$

Eq. (28) is Eq. (3) in section §2.3.1-2 of Polyanin's book, [16]. Therefore, depending on n there are also two sets of parametric solutions, first, if $n\neq-1$ we have

$$\begin{aligned}
&\chi(\tau)=\frac{aC_1^{n-1}}{\Theta(\tau)}\\
&z(\tau)=\frac{bC_1^{n+1}\tau}{\Theta(\tau)}\\
&\Theta(\tau)=C_2+\int\frac{d\tau}{\sqrt{1\pm\tau^{n+1}}}\\
&A=\pm\tfrac{n+1}{2}\ a^{n+1}b^{1-n}\ ,
\end{aligned} \tag{42}$$

while, if $n=-1$, we have

$$\begin{array}{l}\chi(\tau)=\frac{C_1}{\Theta(\tau)}\\ z(\tau)=\frac{b\exp(\mp\tau^2)}{\Theta(\tau)}\\ \Theta(\tau)=C_2+\int\exp(\mp\tau^2)d\tau\\ A=\mp 2b^2\ .\end{array}\tag{43}$$

4 Examples

We present three physical examples, $n=5$ of the first class which occurs in astrophysics of stars treated as gas spheres, while the other ones are $n=2$ cases of the first class and second class. In the latter cases, the equations are particular cases of equations of the type

$$z''=F(\chi)z^2\ ,\qquad n\neq 0\tag{44}$$

which characterize perfect fluids in shear-free motion in general relativity.

In 1948, Kustaanheimo and Qvist found that Eq. (44) is integrable when $F(\chi)=(\alpha_1\chi^2+\alpha_2\chi+\alpha_3)^{-5/2}$ for arbitrary constants α_1, α_2, and α_3 [17]. The case from the first class corresponds to $\alpha_1=\alpha_3=0$, $\alpha_2=(2/3)^{5/2}$, while the case from the second class is the particular case $\alpha_2=\alpha_3=0$, $A=\alpha_1^{-5/2}$.

4.1 $n=2$ of the first class

Let us consider the equation

$$z''+\frac{3}{2}\chi^{-\frac{5}{2}}z^2=0.\tag{45}$$

The first integral of motion is obtained from Eq. (21) using $n=2$, and $A=-\frac{3}{2}$ to obtain

$$z'(\chi z'-z)+\left(\frac{z}{\sqrt{\chi}}\right)^3=\mathcal{C}_1\tag{46}$$

which has two simple rational solutions provided that $\mathcal{C}_1=0$

$$\begin{array}{l}z(\chi)=\frac{4a_1\chi}{(B_2+4\sqrt{\chi})^2}\\ z(\chi)=\frac{4a_1\chi}{(4B_2+\sqrt{\chi})^2},\end{array}\tag{47}$$

with B_2 arbitrary integration constant.

Using the system (41) with $n=2$, we have

$$\begin{array}{l}\chi(\tau)=aB_1^2\exp\Theta(\tau)\\ z(\tau)=bB_1\tau\exp\frac{1}{2}\Theta(\tau)\\ \Theta(\tau)=\int\frac{d\tau}{\sqrt{K_3+\frac{2\tau^3}{3}+\frac{\tau^2}{4}}}\ .\end{array}\tag{48}$$

By inverting, $\tau(\Theta)$ satisfies the elliptic equation

$$\left(\frac{d\tau}{d\Theta}\right)^2=\frac{2\tau^3}{3}+\frac{\tau^2}{4}+K_3=a_3\tau^3+a_2\tau^2+a_1\tau+a_0\equiv Q_3(\tau)\ ,\tag{49}$$

which is put in standard form using the scale-shift transformation

$$\tau(\Theta)=\frac{4}{a_3}\wp(\Theta;g_2,g_3)-\frac{a_2}{3a_3}=6\wp(\Theta;g_2,g_3)-\frac{1}{8}\tag{50}$$

to become

$$\wp_\Theta^2 = 4\wp^3 - g_2\wp - g_3 \ . \tag{51}$$

The germs of the Weierstrass function are given by

$$\begin{array}{l} g_2 = \frac{a_2^2 - 3a_1a_3}{12} = \frac{1}{192} \\ g_3 = \frac{9a_1a_2a_3 - 27a_0a_3^2 - 2a_2^3}{432} = -\frac{1}{13824} - \frac{K_3}{36} \end{array} \tag{52}$$

and together with the modular discriminant [18]

$$\Delta = g_2^3 - 27g_3^2 = -\frac{K_3(1 + 192K_3)}{9216} \tag{53}$$

are used to classify the solutions of Eq. (51).

Case (1). If $\Delta \equiv 0 \Rightarrow K_3 = 0$ or $K_3 = -\frac{1}{192}$ the Weierstrass solutions can be simplified since $\wp$ degenerates into *hyperbolic* or *trigonometric* functions

Case (1a). $K_3 = 0 \Rightarrow g_2 > 0,\ g_3 < 0$ so Eq. (51) has soliton solution

$$\tau(\Theta) = \frac{3}{8}\left[-1 + \tanh^2\left(\frac{\Theta}{4}\right)\right] \tag{54}$$

Case (1b). $K_3 = -\frac{1}{192} \Rightarrow g_2 > 0,\ g_3 > 0$, so Eq. (51) has periodic solution

$$\tau(\Theta) = \frac{1}{8}\left[1 + 3\tan^2\left(\frac{\Theta}{4}\right)\right] \tag{55}$$

Case (2). If $\Delta \neq 0$ the Weierstrass solutions cannot be simplified, but we still have one simpler particular *lemniscatic* case for which $g_2 > 0, g_3 = 0 \Rightarrow K_3 = -\frac{1}{384}$. Then the solution of Eq. (51) is

$$\tau(\Theta) = -\frac{1}{8} + 6\wp\left(\Theta, \frac{1}{192}, 0\right) \tag{56}$$

For all other values of K_3, one obtains the solutions in terms of the general $\wp$ functions. Using the same ansatz as above with particular solution of Eq. (45) given by Eq. (18) we obtain

$$\ddot{\nu} = \frac{1}{4}(\nu - \nu^2) \tag{57}$$

which is the particular case $n = 2$ of Eq. (25). All solutions are presented in Fig. 1.

4.2 $n = 5$ of the first class

In 1907, Emden reviewed the theory of polytropic gas spheres (stars in astrophysics) and studied equations of the form [1]

$$z'' = \pm\chi^{1-n}z^n \ , \tag{58}$$

which belongs to the first class if one chooses $n = 5$. Letting $A = 1$ in Eq. (17), then we obtain

$$z'' = \chi^{-4}z^5. \tag{59}$$

This equation was fist derived by Schuster [19] and has the first integral of motion

$$z'(\chi z' - z) - \frac{1}{3}\left(\frac{z^2}{\chi}\right)^3 = \mathcal{C}_1. \tag{60}$$



Figure 1: Solitonic, periodic, and lemniscatic solutions of the EF equation of the first class with $n = 2$.

Again, if $\mathcal{C}_1 = 0$, the above has solutions

$$z(\chi) = \pm \frac{\sqrt{6} B_3 \chi}{\sqrt{B_3^4 - 12\chi^2}} \tag{61}$$

with B_3 arbitrary integration constant. According to system (41), the parametric solutions are given by

$$\begin{aligned} \chi(\tau) &= aB_1^2 \exp \Theta(\tau) \\ z(\tau) &= bB_1 \tau \exp \tfrac{1}{2}\Theta(\tau) \\ \Theta(\tau) &= \int \frac{d\tau}{\sqrt{K_3 + \frac{\tau^6}{3} + \frac{\tau^2}{4}}} \end{aligned} \tag{62}$$

from which we can see that the solutions can be expressed in terms of Weierstrass elliptic functions.

To find these solutions of Eq. (59), we proceed by using the particular solution of Eq. (59) given by Eq. (18) to obtain

$$\ddot{\nu} + \frac{\nu}{4}(\nu^4 - 1) = 0 \tag{63}$$

which becomes the elliptic equation

$$\dot{u}^2 = -u^4 + u^2 + K_4 u \equiv Q_4(u), \tag{64}$$

using $\nu^2 = \sqrt{3}u$. According to Whittaker and Watson [20] the solutions of Eq. (64) can be expressed in terms of Weierstrass elliptic functions $\wp(t; g_2, g_3)$ via

$$u(t) = u_0 + \frac{\sqrt{Q_4(u_0)}\wp'(t; g_2, g_3) + \frac{1}{2}\frac{dQ_4}{du}(u_0)\Big(\wp(t; g_2, g_3) - \frac{1}{24}\frac{d^2Q_4}{du^2}(u_0)\Big) + \frac{1}{24}Q_4(u_0)\frac{d^3Q_4}{du^3}(u_0)}{2\Big(\wp(t; g_2, g_3) - \frac{1}{24}\frac{d^2Q_4}{du^2}(u_0)\Big)^2 - \frac{1}{48}Q_4(u_0)\frac{d^4Q_4}{du^4}(u_0)}, \tag{65}$$

where u_0 is any constant and not necessarily a root of $Q_4(u)$. The elliptic invariants are related to the coefficients of the quartic polynomial $Q_4(u)$ which yield $g_2 = \frac{1}{12}$, $g_3 = \frac{K_4^2}{16} - \frac{1}{216}$. Because we already know a simple root u_0 of $Q_4(u)$, then Eq. (50) has the simpler form

$$u(t) = u_0 + \frac{\frac{dQ_4}{du}(u_0)}{4\Big(\wp(t; g_2, g_3) - \frac{1}{24}\frac{d^2Q_4}{du^2}(u_0)\Big)}. \tag{66}$$



Figure 2: The periodic and lemniscatic solutions of the EF equation of the first class with $n = 5$.

Choosing $u_0 = 0$ we have the solution

$$u(t) = \frac{K_4}{4\wp\left(t; \frac{1}{12}, \frac{K_4^2}{16} - \frac{1}{216}\right) - \frac{1}{3}}, \qquad K_4 \neq 0. \tag{67}$$

This solution can be reduced depending on the choice of the arbitrary constant K_4, as in the previous case of the similar solution (50) as follows. Using the germs the discriminant is $\Delta = \frac{K_4^2}{256}(4 - 27K_4^2)$. Thus,

Case (1). If $\Delta \equiv 0$, and since $K_4 \neq 0 \Rightarrow K_4 = \pm\frac{2\sqrt{3}}{9}$, therefore $g_3 = \frac{1}{216}$ and the Weierstrass solutions simplifies to the *periodic* solution only

$$u(t) = \pm\frac{\frac{2\sqrt{3}}{3}}{1 + 3\tan^2\left(\frac{t}{2}\right)} \tag{68}$$

Case (2). If $\Delta \neq 0$ the Weierstrass solutions cannot be simplified, but we still have one simpler particular *lemniscatic* case for which $g_3 = 0 \Rightarrow K_4 = \pm\frac{\sqrt{6}}{9}$. Then the solution (67) simplifies to

$$u(t) = \pm\frac{\frac{\sqrt{6}}{3}}{-1 + 12\wp(t; \frac{1}{12}, 0)} \tag{69}$$

Both solutions are presented in Fig. 2.

4.3 $n = 2$ of the second class

We consider now the following equation

$$z'' = A\chi^{-5}z^2 \ . \tag{70}$$

Using the transformation $z(\chi) = \frac{w(s)}{s}$, where $s = \frac{1}{\chi}$, leads to

$$w_{ss} = Aw^2 \tag{71}$$



Figure 3: Solution (76) for the EF equation of the second class with $n = 2$. Notice that this solution cannot be reduced to the equianharmonic case $\wp(\Theta; 0, 1)$ because the reduction requires $g_3 > 0$.

whose integral of motion is

$$w_s^2 + \frac{2}{3}w^3 = \mathcal{C}_3 \ . \tag{72}$$

In terms of the original variables this invariant becomes

$$\chi^4 \left[\frac{d}{d\chi}\left(\frac{z}{\chi}\right)\right]^2 + \frac{2}{3}\left(\frac{z}{\chi}\right)^3 = \mathcal{C}_3 \ . \tag{73}$$

Using $A = -1$ the parametric solutions are

$$\begin{aligned} \chi(\tau) &= \tfrac{aC_1}{\Theta(\tau)} \\ z(\tau) &= \tfrac{bC_1^3\tau}{\Theta(\tau)} \\ \Theta(\tau) &= C_2 + \textstyle\int \tfrac{d\tau}{\sqrt{1\pm\tau^3}}. \end{aligned} \tag{74}$$

By inverting, $\tau(\Theta)$ satisfies the reduced elliptic equation

$$\left(\frac{d\tau}{d\Theta}\right)^2 = \pm\tau^3 + 1 \tag{75}$$

which is a particular case $a_3 = \pm 1$, $a_2 = a_1 = 0$, and $a_0 = 1$ of Eq. (49). The Weierstrass germs are $g_2 = 0$ and $g_3 = -\frac{1}{16}$, therefore $\Delta = -\frac{3^3}{2^8}$ and the solution to Eq. (75) is given by

$$\tau(\Theta) = \pm 4\wp\left(\Theta; 0, -\frac{1}{16}\right). \tag{76}$$

A plot of this solution for the positive sign is given in Fig. 3. Using the particular solution of Eq. (70) given by Eq. (29) in Eq. (16), we obtain

$$\ddot{\nu} + 5\dot{\nu} + 6(\nu - \nu^2) = 0 \tag{77}$$

and using the factorization technique becomes

$$\dot{\nu} + 2(\nu - \nu^{\frac{3}{2}}) = 0 \tag{78}$$

with solution

$$\nu(t) = \left(\frac{1}{1 + K_2 e^t}\right)^2 , \tag{79}$$

as obtained from Eq. (39).

5 Conclusion

In summary, we have provided a discussion of the two integrable classes of Emden-Fowler equations, $\lambda = \frac{n-1}{2}$ and $\lambda = n + 1$. As examples, in the first class, we have presented the astrophysical case $\lambda = 2$, $n = 5$ which lead to Weierstrass elliptic solutions, and two $n = 2$ cases, one with $\lambda = 1/2$ belonging to the first class and another one with $\lambda = 3$ from the second class. The $n = 2$ cases usually describe perfect fluids in general relativity. Many other cases with potential applications, such as those with negative n, which belong to these two integrable classes can be approached along the lines presented in this work.

References

[1] R. Emden, Gas balls: Applications of the Mechanical Heat Theory to Cosmological and Meteorological Problems, Teubner, (1907) (in German).

[2] J.H. Lane, On the theoretical temperature of the sun under the hypothesis of a gaseous mass maintaining its volume by its internal heat and depending on the laws of gases known to terrestrial experiment, Am. J. Sci. 50, (1870) 57-74.

[3] Sir A.S. Eddington, The Internal Constitution of the Stars, Cambridge Univ. Press, (1926).

[4] R.H. Fowler, The solutions of Emden's and similar differential equations, Monthly Notices of the Royal Astronomical Society 91 (1930) 63-91.

[5] E.A. Milne, The analysis of stellar structure, Monthly Notices of the Royal Astronomical Society 91 (1930) 4-52.

[6] A.W. Rodgers, D.M. Myers, Solutions of the negative Emden polytropes, Monthly Notices of the Royal Astronomical Society 114 (1954) 620-627.

[7] J. Janus, J. Myjak, A generalized EF equation with a negative exponent, Nonlinear Analysis: Theory, Methods & Applications 23, (1994) 953-970.

[8] L.H. Thomas, The calculation of atomic fields, Proc. Cambridge Phil. Soc. 23, (1927) 542-548.

[9] E. Fermi, A statistical method for the determination of some atomic properties, Rend. Accad. Naz. Lincei 6 (1927) 602-607 (in Italian).

[10] R.H. Fowler, The form near infinity of real continuous solutions of a certain differential equation of the second order, Quart. J. Math. (Oxford) 45, (1914) 289.

[11] E. Kamke, Differentialgleichungen. Lösungsmethoden und Lösungen I. Gewöhnliche Differentialgleichungen, Chelsea Publ. Co., 1948, eqs. 6.11 at p. 544 and 6.74 at p. 560.

[12] P. Rosenau, A note on the integration of the Emden-Fowler equation, Int. J. Non-Linear Mech. 19, (1984) 303-308.

[13] D.S. Djukic, A procedure for finding first integrals of mechanical systems with gauge-variant Lagrangians, Int. J. Non-Linear Mech. 8, (1973) 479-488.

[14] R.H. Fowler, Further studies of Emden's and similar differential equations, Quart. J. Math. (Oxford) 2, (1931) 259-288.

[15] H.C. Rosu, O. Cornejo-Pérez, Supersymmetric pairing of kinks for polynomial nonlinearities, Phys. Rev. E 71, (2005) 046607.

[16] A.D. Polyanin, V.F. Zaitsev, Handbook of exact solutions for ordinary differential equations, CRC Press, (1995).

[17] P. Kustaanheimo, B. Qvist, A note on some general solutions of the Einstein field equations in a spherically symmetric world, Societas Sci. Fennica. Comm. Physico-Mathematicae XIII.16, 1 (1948). Reprinted in Gen. Rel. Grav. 30, (1998) 663-673.

[18] M. Abramowitz, I. Stegun, Handbook of Mathematical Functions: with Formulas, Graphs, and Mathematical Tables, Dover, N.Y., (1972)

[19] A. Schuster, On the internal constitution of the Sun, Report of the British Association for the Advancement of Science (1883), 427-429.

[20] E.T. Whittaker, G.N. Watson, A Course of Modern Analysis, Cambridge University Press, Cambridge, (1927).